

The American Journal of Sports Medicine

<http://ajs.sagepub.com/>

Ultrasound-Guided Sclerosing Treatment in Patients With Patellar Tendinopathy (Jumper's Knee) : 44-Month Follow-up

Aasne Hoksrud and Roald Bahr

Am J Sports Med 2011 39: 2377 originally published online August 12, 2011

DOI: 10.1177/0363546511417097

The online version of this article can be found at:

<http://ajs.sagepub.com/content/39/11/2377>

Published by:



<http://www.sagepublications.com>

On behalf of:



American Orthopaedic Society for Sports Medicine

Additional services and information for *The American Journal of Sports Medicine* can be found at:

Email Alerts: <http://ajs.sagepub.com/cgi/alerts>

Subscriptions: <http://ajs.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [Version of Record](#) - Oct 28, 2011

[Proof](#) - Aug 12, 2011

[What is This?](#)

Ultrasound-Guided Sclerosing Treatment in Patients With Patellar Tendinopathy (Jumper's Knee)

44-Month Follow-up

Aasne Hoksrud,^{*†} MD, and Roald Bahr,[†] MD, PhD

Investigation performed at the Oslo Sports Trauma Research Center, Norwegian School of Sport Sciences, Oslo, Norway

Background: A randomized controlled study has shown good clinical results after treatment with sclerosing injections into the area with neovessels in patients with patellar tendinopathy, but no study has investigated medium- or long-term outcomes.

Purpose: This study investigates the effect of sclerosing treatment 44 months (range, 42-47 months) after start of treatment.

Study Design: Case series; Level of evidence, 4.

Methods: Patients with a diagnosis of jumper's knee and neovascularization corresponding to the painful area were recruited and treated with ultrasound-guided sclerosing injections using polidocanol. Primary outcome was Victorian Institute of Sport Assessment (VISA) score, which was recorded before the start of treatment, after 12 months, and 44 months after the start of the study period.

Results: Twelve of the 29 patients (14 tendons) who were followed up at 44 months had undergone arthroscopic surgery after sclerosing treatment, either to the patellar tendon ($n = 6$) or for other intra-articular lesions ($n = 8$). For patients who did not receive additional treatment after the sclerosing injections ($n = 23$ tendons), VISA score was 55 (range, 28-71) at baseline and 81 (range, 39-100) at 12-month follow-up ($P < .001$ vs baseline). Their VISA score at 44 months' follow-up was 89 (range, 73-100; $P = .047$ vs 12 months). For patients who went through arthroscopic tendon surgery, VISA score was 53 (range, 39-71) at baseline and 71 before surgery (range, 48-98; $P = .14$ vs baseline). Their VISA score at 44 months was 91 (range, 76-100; $P = .016$ vs 12 months; $P = .005$ vs baseline). For patients who went through non-tendon surgery, VISA score was 45 (range, 15-69) at baseline and 57 (range, 32-95) before surgery ($P = .29$ vs baseline). Their VISA score at 44 months was 92 (range, 72-100; $P = .006$ vs before surgery; $P < .001$ vs baseline).

Conclusion: Sclerosing treatment with polidocanol was effective for the majority of the patients. Nevertheless, one-third elected to seek additional treatment through arthroscopic surgery during the 44-month follow-up period.

Keywords: jumper's knee; polidocanol; color Doppler; tendon

The exact causes of pain in tendinopathy are unclear, but may be related to increased vascularity in the tendon.^{2,12} Studies have shown that neovascularization is present in

60% to 80% of patients with chronic painful patellar tendinopathy,⁵ and the presence of neovascularization was associated with more tendon pain than in abnormal tendons without neovascularization.^{5,6} Based on the hypothesis that these vessels and their accompanying nerves are involved in the pain mechanism, a pilot study on sclerosing treatment for patients with Achilles midportion tendinopathy reported promising results.¹¹ This led to a randomized clinical trial investigating the effect of polidocanol injections in 20 patients with Achilles tendinopathy, demonstrating a significant short-term improvement compared with placebo injections.¹ In a study investigating the 2-year outcome after sclerosing treatment in Achilles tendinopathy, 38 of 42 patients reported treatment satisfaction.¹⁰

On the basis of the promising results in Achilles tendinopathy, we completed a randomized controlled trial of polidocanol injection therapy in elite athletes with patellar tendinopathy, where the patients were randomized to either immediate or delayed polidocanol injections.⁷ The 12-month

*Address correspondence to Aasne Fenne Hoksrud, MD, Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, PO Box 4014 Ullevaal Stadion, 0806 Oslo, Norway (e-mail: aasne.hoksrud@nih.no).

[†]Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway.

One of more of the authors has declared the following potential conflict of interest or source of funding: The Oslo Sports Trauma Research Center has been established at the Norwegian School of Sport Sciences through generous grants from the Royal Norwegian Ministry of Culture, the South-Eastern Norway Regional Health Authority, the International Olympic Committee, the Norwegian Olympic Committee & Confederation of Sport, and Norsk Tipping AS.

results have been described in detail and led us to conclude that knee function and pain improved significantly.⁷ However, no study has investigated the medium- or long-term outcome after polidocanol treatment in patients with patellar tendinopathy. Therefore, in this article, we report on the activity levels, knee function, and overall treatment satisfaction of the patients included in the randomized trial after 44 months.

METHODS

The randomized trial included 33 elite athletes (5 female and 28 male athletes) with a mean age of 25 years (range, 17-42 years), mainly representing team handball ($n = 15$), basketball ($n = 5$), and football (soccer) ($n = 6$), with 43 tendons with neovascularization. The team sport athletes competed in the top 2 divisions of the respective Norwegian league, and in the individual sports, the athletes represented a comparable level. The study protocol has been described in detail by Hoksrud et al.⁷

Athletes were invited to a clinical screening examination, and the following diagnostic criteria were used to identify patients with jumper's knee⁸: history of pain in the patellar tendon or the patellar insertion in connection with training or competition, tenderness to palpation corresponding to the painful area,⁴ and symptoms from the patellar tendon for a minimum of 3 months. Patients who fulfilled the diagnostic criteria were invited to an ultrasound examination, including a color Doppler examination to assess neovascularization. To be included in the study, patients had to have a clinical diagnosis of jumper's knee, structural tendon changes and neovascularization corresponding to the painful area on the ultrasound examination, and a Victorian Institute of Sport Assessment (VISA) score (0-100 points) of less than 75 points.

The sclerosing treatment involved ultrasound-guided injections with the sclerosing agent polidocanol (Aethoxysklerol [10 mg/mL], Inverdia AB, Stockholm, Sweden). The patients were scheduled for follow-up visits in the laboratory 3 to 5 weeks after each treatment and those who reported pain and reduced function were offered a new sclerosing injection if they had persistent neovascularization; no further injections were given if no vessels were seen on the ultrasound examination. During the treatment period (0-8 months in the group who received immediate injections, $n = 16$; 4-8 months in the group receiving delayed treatment, $n = 17$), patients received 0 to 5 sclerosing injections (mean, 3.1) with 3- to 5-week intervals (0 injections, 4 tendons; 1 injection, 6 tendons; 2 injections, 10 tendons; 3 injections, 10 tendons; 4 injections, 2 tendons; 5 injections, 11 tendons).

The primary outcome measure was knee function using the VISA score.¹⁴ Secondary outcome was overall satisfaction with treatment using a visual analog scale. For overall satisfaction, we asked the patients to evaluate the sclerosing treatment on a scale from 0 to 10, where 0 represented very unsatisfied with treatment and 10 represented very satisfied with treatment. The outcomes were recorded at baseline, and 12 and 44 months after the first injection. At baseline and the 12-month follow-up, the outcomes

were recorded in writing and at 44 months by a telephone interview.

The study has been approved by Norwegian School of Sport Sciences. Within-group differences were assessed with paired *t* tests using a significance level of 5%, and the results are presented as the mean with their range or 95% confidence interval, as noted.

RESULTS

Of the 33 patients (43 tendons) who were included in the randomized controlled trial and initially followed for 12 months, we were able to follow-up with 29 patients (37 tendons) at 44 months (range, 42-47). The VISA score for the 4 patients (6 tendons) who did not participate in the final follow-up was excellent at 12 months (93; range, 87-98).

Twelve of the 29 patients (14 tendons) who were followed up at 44 months had undergone arthroscopic surgery in the meantime, either to the patellar tendon or for other intra-articular injury. In 6 cases (6 patients), tendon surgery was performed; in 7 cases (6 patients), debridement of minor retropatellar chondral defects was performed; and in 1 case, a plica medialis was resected. Of these, there were 2 patients with bilateral problems; 1 had non-tendon surgery performed on both knees, the other had tendon surgery on 1 knee and non-tendon surgery on the other knee.

In all cases that went through tendon surgery, this was performed after the 12-month follow-up. In the 8 cases that went through non-tendon surgery, 5 surgeries were performed before the 12-month follow-up and 3 after the 12-month follow-up.

Of the 37 tendons that were followed up at 44 months, 24 cases were training and competing at the same level as before injury, while the activity level was reduced in 13 cases, 3 because of knee problems (Figure 1).

VISA Score

The patients who did not receive any additional treatment after the sclerosing injections ($n = 23$ tendons) reported significantly improved VISA scores from baseline (55; range, 28-75) to the 12-month follow-up (81; range, 39-100) ($P < .001$), and a further improvement in pain and function scores from the 12- to the 44-month follow-up (89; range, 73-100; $P = .047$) (Figure 2).

Patients who went through arthroscopic surgery to the tendon (patellar tendon debridement, $n = 6$ tendons) did not report a significant improvement in VISA score from baseline (53; range, 39-71) until before surgery (ie, at 12 months) (71; range, 48-98; $P = .14$). Their VISA score at 44 months was 91 (range, 76-100; $P = .016$ vs 12 months; $P = .005$ vs baseline) (Figure 2).

For patients who went through non-tendon surgery ($n = 8$ tendons), their VISA score was 45 (range, 15-69) at baseline and 57 (range, 32-95) before surgery (ie, at 4 or 12 months; $P = .29$ vs baseline). They reported a significantly improved VISA score at the 44-month follow-up (92; range, 72-100) compared with baseline ($P < .001$) and with their last score before surgery ($P = .006$) (Figure 2).




Figure 1. Flowchart depicting treatments, follow-up, and activity level.

Overall Treatment Satisfaction

Patients who did not receive other treatment after the sclerosing injection were equally satisfied with treatment at 12-month (7.9; range, 3-10) and 44-month follow-up (8.2; range, 4-10; $P = .30$, paired t test). Patients who went through subsequent tendon surgery were more satisfied with treatment before surgery (ie, at 12 months) (7.8; range, 4-10) compared with the 44-month follow-up (3.3; range, 2-5; $P = .001$ vs 12 months). Overall treatment satisfaction for patients who went through subsequent non-tendon surgery was 4.0 (range, 1-8) before surgery (ie, at 4 or 12 months) and 5.8 (range, 3-10) at 44 months ($P = .13$).

DISCUSSION

This follow-up study shows that injection treatment with polidocanol resulted in a significant improvement in knee function and reduced pain 44 months after the start of treatment in patients with patellar tendinopathy. However, it should be noted that in more than one-third of the cases, patients still elected to seek additional treatment through arthroscopic surgery during the follow-up period, either to the tendon itself (16%) or for other intra-articular lesions (22%). These results therefore show that, despite the group improvements documented, a significant proportion of patients were not fully satisfied. The long-term follow-up also shows that there was a subgroup of patients who turned out to have additional intra-articular injuries that may have contributed to their




Figure 2. Victorian Institute of Sport Assessment (VISA) score (mean, 95% confidence interval) for patients who did not receive additional treatment after sclerosing injections, patients who went through tendon surgery, and patients who went through non-tendon surgery at baseline, 12 months, and 44 months. For patients who went through non-tendon surgery, the 12-month follow-up was before surgery (ie, at 4 or 12 months).

symptoms, even though a careful inclusion protocol that included ultrasound imaging was used.

It is perhaps not surprising that many of the patients sought out other treatment options. The average VISA score at 12-month follow-up for all patients included in the randomized controlled trial was 77, which represented a significant improvement from their baseline score of 54.⁷ Nevertheless, the scores ranged widely, from 32 to 100 at 12 months. The current trial is unique in that all patients were elite athletes, and it cannot be expected that they would be fully satisfied with a score of less than 90. However, in as many as 8 cases, the 12-month VISA score was <50 and in 12 cases <60.

The pioneering work on sclerosing treatment was done on Achilles tendinopathy and the majority of patients included in clinical studies were recreational runners, typically aged 43 to 74 years¹¹; the treatment satisfaction and return-to-sport rates they report are not necessarily valid in younger, elite athletes. Return to sport is not a reliable outcome in elite athletes, as they tend to continue to train and compete despite significant pain and loss of function. This can be seen from the cross-sectional study by Lian et al,⁹ which demonstrated that in 87 elite athletes with jumper's knee, who were competing at the national elite level in 9 different sports, the average symptom duration was 32 months and the average VISA score was 64. This can also be seen in the present study, where only 3 patients had retired from elite sports because of knee problems.

In other studies investigating the effects of sclerosing treatment in patients with tendinopathy, interpretation is also confounded by patients undergoing other treatment modalities, such as surgery, before final assessment. Although the results appear to be good, some of the tendons had undergone additional treatment at some point and

treatment contamination may have affected the final outcome. One example is van Sterkenburg et al,¹³ who retrospectively assessed the effects of sclerosing treatment on Achilles tendinopathy after 2.7 to 5.1 years follow-up. Although the results show that many patients improved, as many as 21 of the 40 tendons they were able to follow had undergone additional treatment at some point; 15 tendons of these were treated operatively (open surgical debridement or Achilles tendoscopy).

In the present study, we wanted to prevent confounding by separating the patients into different groups, depending on whether they received additional surgical treatment or not. We have therefore divided the patients into 3 groups: 1 group that did not receive surgical treatment, 1 group of patients who went through tendon surgery, and 1 group that went through non-tendon surgery. All patients taken together, VISA scores improved, but patients who went through surgery did not report a significant improvement in VISA scores from baseline to before surgery. Their VISA scores likely explain why they elected to seek additional treatment. An interesting observation is that patients who went through tendon surgery were less satisfied with sclerosing treatment after surgery (ie, 44 months) compared with before surgery (ie, 12 months), even if their knee function (VISA score) had improved significantly. The most likely explanation is that they attribute their improved function to surgical treatment and, consequently, were less satisfied with the sclerosing treatment than before.

The group of patients who went through non-tendon surgery also illustrates the difficulty with the diagnostic criteria for patellar tendinopathy.^{3,4} We included patients based on a careful history and clinical examination and included an ultrasound examination requiring structural changes and neovascularization. Despite this, in 8 cases, coexisting intra-articular conditions were revealed through subsequent arthroscopic surgery.

The current data also illustrate the difficulty of conducting controlled studies on elite athletes. Ideally, we would like to have long-term data from randomized trials to inform clinical decisions. In the present trial, we were able to randomize patients into immediate treatment or to a group receiving placebo injections during an initial 3-month period. However, it seems highly unlikely that elite athletes would be willing to accept placebo treatment for a sufficient period. For this reason, we may have to continue basing clinical decisions for tendinopathy on short-term outcomes or data from recreational athletes. When interpreting the data from the present study, it should also be borne in mind that follow-up at 44 months was done through a telephone interview, and that the sample size was limited.

In conclusion, this follow-up study shows that sclerosing treatment results in a midterm improvement in knee function and pain in a group of young, elite athletes with patellar tendinopathy. However, few experience complete resolution of symptoms and as many as one-third elected to seek additional treatment through arthroscopic surgery during the 44-month follow-up period.

ACKNOWLEDGMENT

The authors thank Håkan Alfredson, MD, PhD, and Lars Öhberg, MD, PhD, for their collaboration in completing the original randomized clinical trial, on which this follow-up study is based. The authors also thank the athletes who participated in the study.

REFERENCES

1. Alfredson H, Ohberg L. Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc.* 2005;13:338-344.
2. Alfredson H, Ohberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc.* 2003;11:334-338.
3. Cook JL, Khan KM, Kiss ZS, Griffiths L. Patellar tendinopathy in junior basketball players: a controlled clinical and ultrasonographic study of 268 patellar tendons in players aged 14-18 years. *Scand J Med Sci Sports.* 2000;10:216-220.
4. Cook JL, Khan KM, Kiss ZS, Purdam CR, Griffiths L. Reproducibility and clinical utility of tendon palpation to detect patellar tendinopathy in young basketball players: Victorian Institute of Sport tendon study group. *Br J Sports Med.* 2001;35:65-69.
5. Cook JL, Kiss ZS, Khan KM, Purdam CR, Webster KE. Anthropometry, physical performance, and ultrasound patellar tendon abnormality in elite junior basketball players: a cross-sectional study. *Br J Sports Med.* 2004;38:206-209.
6. Hoksrud A, Ohberg L, Alfredson H, Bahr R. Color Doppler ultrasound findings in patellar tendinopathy (jumper's knee). *Am J Sports Med.* 2008;36:1813-1820.
7. Hoksrud A, Ohberg L, Alfredson H, Bahr R. Ultrasound-guided sclerosis of neovessels in painful chronic patellar tendinopathy: a randomized controlled trial. *Am J Sports Med.* 2006;34:1738-1746.
8. Lian O, Holen KJ, Engebretsen L, Bahr R. Relationship between symptoms of jumper's knee and the ultrasound characteristics of the patellar tendon among high level male volleyball players. *Scand J Med Sci Sports.* 1996;6:291-296.
9. Lian OB, Engebretsen L, Bahr R. Prevalence of jumper's knee among elite athletes from different sports: a cross-sectional study. *Am J Sports Med.* 2005;33:561-567.
10. Lind B, Ohberg L, Alfredson H. Sclerosing polidocanol injections in mid-portion Achilles tendinosis: remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2006;14:1327-1332.
11. Ohberg L, Alfredson H. Ultrasound guided sclerosis of neovessels in painful chronic Achilles tendinosis: pilot study of a new treatment. *Br J Sports Med.* 2002;36:173-175.
12. Ohberg L, Lorentzon R, Alfredson H. Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: an ultrasonographic investigation. *Knee Surg Sports Traumatol Arthrosc.* 2001;9:233-238.
13. van Sterkenburg MN, de Jonge MC, Siersevelt IN, van Dijk CN. Less promising results with sclerosing ethoxysclerol injections for mid-portion Achilles tendinopathy: a retrospective study. *Am J Sports Med.* 2010;38:2226-2232.
14. Visentini PJ, Khan KM, Cook JL, Kiss ZS, Harcourt PR, Wark JD. The VISA score: an index of severity of symptoms in patients with jumper's knee (patellar tendinosis)—Victorian Institute of Sport Tendon Study Group. *J Sci Med Sport.* 1998;1:22-28.